

## Qualitative differences between naïve and memory T cells

MARION BERARD & DAVID F. TOUGH *The Edward Jenner Institute for Vaccine Research, Compton, Newbury, Berkshire RG20 7NN, UK*

### INTRODUCTION

Mature T cells are produced in the thymus and released into the bloodstream in low numbers. These cells are considered to be immunologically naïve until such time as they encounter MHC-peptide complexes for which their T-cell receptors (TCR) have high affinity. Recognition of antigen in appropriate form, i.e. in association with costimulatory signals on the surface of professional antigen-presenting cells (APCs), leads to extensive T-cell proliferation and differentiation into effector cells. Once the infection has been cleared, it is no longer of benefit to the host to maintain high numbers of effector cells and most of the activated T cells die by apoptosis. However, a proportion of these cells survive, leaving the frequency of cells specific for the priming antigen much higher among memory T cells than that which existed among naïve T cells. This difference in frequency makes a major contribution to the nature of the secondary response, which is typically faster and of greater magnitude than the primary response. In addition, T cells may also carry a true 'memory' of a prior response to antigen, exhibiting differences from naïve T cells at the single cell level. Here we provide a brief overview of the qualitative differences that have been reported to exist between naïve and memory T cells and evidence that memory T cells themselves are functionally heterogeneous.

### PHENOTYPIC DIFFERENCES BETWEEN NAÏVE AND MEMORY T CELLS

The supposition that naïve and memory T cells can be distinguished phenotypically is based on the notion that memory T cells retain a permanent imprint of having responded to antigen. Precise identification of memory T cells, however, remains problematic. Unlike B cells, T cells do not appear to mutate their antigen receptor genes during the course of an immune response. Furthermore, discrimination between effector and memory T cells is accomplished

on the basis of rather nebulous criteria; memory T cells are considered to differ from effector T cells by their continued survival after the acute immune response has died down and by being in a lower state of activation. As discussed further below, these distinctions are becoming increasingly blurred.

Despite these difficulties, a number of phenotypic differences between naïve and memory T cells have been noted. Most of these are changes that arise during initial T-cell activation and appear to persist in memory cells. Especially prominent are differences in the cell surface expression of adhesion molecules between naïve and memory T cells. Thus, compared to naïve T cells, memory T cells have been reported to express higher levels of  $\beta 1$  (CD29, CD49d and CD49e) and  $\beta 2$  (CD11a, CD11b and CD18) integrins, CD2, CD44, CD54 and CD58.<sup>1–11</sup> Increased expression of adhesion molecules on recently activated T cells reflects the requirements for effector T cells to enter peripheral tissues at sites of inflammation and interact with target cells, and may similarly affect the function of some memory T cells (see below).

The expression of other molecules involved in lymphocyte migration also differs between naïve and memory T cells. Of particular interest are differences in the expression of two key molecules required for the entry of T cells into lymph nodes through high endothelial venules (HEVs): CD62L and CCR7. CD62L binds to vascular addressins expressed on HEVs and is responsible for the initial stage of adherence of blood-borne T cells to HEVs,<sup>12</sup> while the CCR7 chemokine receptor controls responsiveness to chemokines expressed in HEVs at sites of lymphocyte entry.<sup>13</sup> Whereas naïve T cells are uniform in expressing high levels of both molecules, some memory cells lose expression of CD62L and/or CCR7.<sup>14–17</sup> However, memory T cells may express receptors for chemokines that direct them to inflammatory sites and for molecules involved in homing to peripheral tissues, such as the cutaneous lymphocyte antigen (CLA) which is involved in lymphocyte migration to skin.<sup>17</sup>

Other cell surface molecules that have been reported to distinguish between naïve and memory T cells include the IL-2R  $\beta$ -chain (CD122),<sup>18,19</sup> Ly-6C<sup>19–21</sup> and the common leukocyte antigen (CD45).<sup>4,6,14,22–28</sup> CD122 is a component of both the IL-2R and the IL-15R and may play a role in the maintenance of memory T cells (see below),<sup>29</sup> while

Received 2 April 2002; accepted 17 April 2002.

Correspondence: David F. Tough, The Edward Jenner Institute for Vaccine Research, Compton, Newbury, Berkshire RG20 7NN, UK. E-mail: david.tough@jenner.ac.uk

Ly-6C is a low-molecular-weight (MW) glycosylphosphatidylinositol-anchored molecule that has been proposed to participate in intercellular adhesion;<sup>30</sup> both CD122 and Ly-6C are expressed at high levels on CD8<sup>+</sup> memory T cells in the mouse. For CD45, which is a tyrosine phosphatase that regulates signalling through antigen receptors and cytokine receptors,<sup>31,32</sup> it is the form of the molecule that differs between naïve and memory cells, rather than the level of expression. Multiple isoforms of CD45 are generated by differential splicing of three extracellular exons (A, B and C). These restricted (R) isoforms can be detected specifically with mAb directed against the variably spliced exons. Naïve T cells express the highest MW isoform, containing all three of these exons (commonly referred to as CD45RA in humans). During the course of T-cell activation, T cells switch to expressing lower-MW isoforms; in humans at least, activated T cells express the isoform of CD45 lacking all three variably spliced exons (defined as CD45R0). In many different species, expression of low-MW isoforms of CD45 is retained on memory cells.

Since many of the phenotypic properties associated with memory T cells are in fact acquired soon after activation, these markers cannot be used on their own to discriminate between recently activated cells and memory cells. This distinction can be aided to a certain extent by combining these phenotypic markers with other criteria to exclude T cells that are actively responding to antigen. For example, it is generally assumed that memory T cells do not have a blasted morphology and do not express transient markers of activation such as CD69. Furthermore, differences in cell surface glycosylation have been reported to exist between effector and memory CD8<sup>+</sup> T cells in mice. Specifically, memory cells have a higher degree of sialylation on 1 O-glycans and express lower levels of 2 O-glycans than effector cells;<sup>33–36</sup> this difference can be detected using an antibody that binds specifically to CD43 only when this molecule has been modified by 2 O-glycans.<sup>37</sup> However, in using signs of recent activation as exclusion criteria for memory T cells, it must be borne in mind that even long-term memory cells appear to be more metabolically active than naïve T cells.<sup>38</sup> As discussed below, memory T cells undergo periodic rounds of cell division even in the complete absence of antigen. Therefore, markers of recent activation cannot be used definitively to distinguish between effector and memory T cells.

Although memory T cells are enriched amongst cells expressing the surface markers discussed above, it is also clear that memory cells exhibit substantial phenotypic heterogeneity. This issue has received considerable attention in recent years, with interest stemming largely from a report that CD45R0<sup>+</sup> T cells in human blood can be divided into CD62L<sup>+</sup>CCR7<sup>+</sup> and CD62L<sup>−</sup>CCR7<sup>−</sup> subpopulations.<sup>17</sup> These cells have been termed 'central memory' and 'effector memory' cells, respectively, based on their expression of lymph node homing molecules and their functional properties (see below). In addition to CD62L and CCR7, memory T cells may express other markers associated with naïve T cells, such as high-MW isoforms of CD45.<sup>16,17,39–46</sup>

In some instances, it is evident that expression of a 'naïve' phenotype by primed cells represents phenotypic reversion. This has been shown to be the case for CD62L, CCR7 and high-MW isoforms of CD45, each of which can be re-expressed by cells that were formerly negative for these markers.<sup>39,41,47–52</sup> Phenotypic reversion occurs at different rates in different species and also differs for CD4 vs. CD8 cells. For example, rat CD45RC<sup>−</sup> (memory-phenotype) CD4<sup>+</sup> T cells re-express CD45RC within 1 week when transferred to secondary recipients in the absence of antigen;<sup>39,41</sup> the rapidity with which this reversion takes place suggests that the CD45RC<sup>−</sup> phenotype in the rat is a marker of recent activation rather than memory. Conversely, CD4<sup>+</sup> memory T cells in mice can maintain a CD45RB<sup>low</sup> phenotype for at least 10 weeks in the absence of antigen, although CD8<sup>+</sup> T cells re-express CD45RB soon after activation.<sup>53</sup>

Phenotypic reversion is presumed to reflect a 'cooling down' of activated cells; lack of contact with antigen results in a return to a resting state and the loss of expression of activation molecules. By corollary, retention of memory markers may be indicative of periodic contact with persisting antigen. However, some phenotypic markers, particularly the expression of high levels of CD44 on mouse memory T cells, appear to be retained long-term in the complete absence of antigen.<sup>54,55</sup> In addition, mechanisms other than reversion may account for some of the phenotypic heterogeneity observed amongst memory T cells. In this respect, it is notable that some CD45RA<sup>+</sup> CD8<sup>+</sup> T cells in human peripheral blood exhibit the properties of activated effector cells.<sup>11</sup> These cells, which also express low levels of CD28 and CD27, appear to arise from chronic antigenic stimulation.<sup>56,57</sup> Likewise, some primed CD45RA<sup>+</sup> CD4<sup>+</sup> T cells can be found under conditions of chronic antigen exposure.<sup>45</sup>

Whether the CD45RA<sup>+</sup> cells observed in these studies have in fact re-expressed this molecule is unclear. Another possibility is that some cells may retain expression of CD45RA under certain conditions of activation. This is worth considering in view of data showing that mouse CD8<sup>+</sup> T cells can differentiate directly into cells with the properties of central memory cells following antigenic stimulation *in vitro*.<sup>58,59</sup> In these studies, brief exposure of CD8<sup>+</sup> T cells to antigen followed by culture in IL-15 or low doses of IL-2 generated T cells that retained expression of CD62L and CCR7 and which lacked overt effector activity. By contrast, cells exposed to high concentrations of IL-2 after antigenic stimulation lost expression of CD62L and CCR7 and differentiated into effector cells. Therefore, acquisition of all of the markers typically associated with T-cell activation is not an inevitable consequence of antigenic stimulation.

Also complicating the use of phenotypic markers to distinguish between naïve and memory T cells is the fact that naïve T cells can acquire markers of memory cells in the absence of overt antigenic stimulation. This has been demonstrated to occur when small numbers of naïve cells are adoptively transferred into lymphopenic recipients.<sup>60–68</sup> Under these conditions, naïve T cells proliferate slowly

(so-called homeostatic proliferation), up-regulate expression of activation/memory markers and exhibit effector activity. Notably, this response is driven not by specific antigen but by self-peptide-MHC complexes in combination with cytokines.<sup>60,61,64,69–71</sup> The contribution of cells generated by this process to the pool of memory-phenotype T cells in normal mice is unclear. However, the fact that very few memory-phenotype T cells are observed in germ-free mice argues that most memory-phenotype T cells are derived from antigenic stimulation.<sup>27</sup> Nevertheless, it remains possible that the phenotypic conversion associated with homeostatic proliferation may make a significant contribution to the pool of memory-phenotype T cells under conditions of lymphopenia.

### NAÏVE AND MEMORY T CELLS EXHIBIT QUALITATIVELY DIFFERENT RESPONSES TO ANTIGEN

It has been evident for many years that memory-phenotype T cells respond to antigen in a qualitatively different way from naïve-phenotype T cells.<sup>2,27,72–81</sup> In recent years, experiments employing TCR transgenic T cells have provided strong evidence that this is also true for bona fide naïve and memory cells. Differences are manifest at two levels.

Firstly, memory T cells appear to have less stringent requirements for activation than naïve T cells. This may include an ability to respond to lower concentrations of antigen than naïve T cells,<sup>21,82–85</sup> although some investigators have failed to find any difference between naïve and memory T cells in their sensitivity to antigen.<sup>86–88</sup> In addition, memory T cells are less dependent on costimulatory signals than naïve T cells, and do not require as long a duration of antigenic stimulation. Each of these factors may contribute to the fact that a wider range of cells can act as APCs for memory T cells compared to naïve T cells. Thus, while activation of naïve T cells is strictly dependent on antigen presentation by dendritic cells (DCs),<sup>89</sup> memory T cells respond to antigen presented on other APCs, including resting B cells.<sup>90</sup>

Secondly, once they have been activated, the response characteristics of naïve and memory T cells also differ. For example, there is some evidence that memory T cells proliferate faster and reach higher numbers *in vivo* than naïve T cells following antigenic stimulation.<sup>91–93</sup> However, this issue remains contentious, as these differences have not been detected in other studies.<sup>88</sup> Conversely, it is a general finding that memory T cells display effector functions sooner after activation than naïve T cells. This includes the expression of cytolytic activity (for CD8<sup>+</sup> T cells) and the secretion of cytokines other than IL-2.

Some clues as to the biochemical basis of the altered responsiveness of memory T cells have been uncovered. For instance, the rapid expression of effector cytokines by memory T cells may be linked to the modulation of chromatin structure or demethylation of promoters for cytokine genes; this has been shown to occur following

initial T-cell activation and is inherited by daughter cells.<sup>94,95</sup> Indeed, some memory T cells have been shown to express mRNA for effector cytokines prior to secondary stimulation.<sup>92,93</sup> In addition, non-dividing memory T cells possess a higher content of both RNA and protein than naïve T cells, suggesting that memory cells may be resting in the G<sub>1</sub> rather than the G<sub>0</sub> phase of the cell cycle.<sup>38,93</sup> This might allow for faster entry into S phase and initiation of DNA replication by memory T cells following TCR triggering. Furthermore, several differences between naïve and memory T cells have been noted that could affect signalling pathways within these cells. Specifically, naïve and memory T cells have been shown to differ with respect to phosphorylation and association of signalling components of the TCR/CD3 complex,<sup>96</sup> expression of the linker/adaptor molecule SLP-76<sup>97</sup> and association of CD45 with the TCR or CD4.<sup>98,99</sup>

Another factor that could contribute to the altered properties of memory T cells is an enhanced ability to interact with APCs compared to naïve T cells. Given their increased expression of various adhesion molecules (see above), it seems likely that memory cells will form higher avidity interactions with other cells; this could contribute to the wider range of APCs utilized by memory vs. naïve T cells. Furthermore, there is evidence that T cells are selectively recruited on the basis of expressing TCRs with higher affinity for antigen during the course of an immune response.<sup>100–110</sup> This is a result of the loss of cells expressing TCRs with the fastest dissociation rates for peptide-MHC binding.<sup>104,106</sup> Since TCR genes do not undergo somatic hypermutation, this process is not directly analogous to the affinity maturation involved in the generation of memory B cells, and results in relatively modest increases in affinity. Nevertheless, the data suggest that memory T cells may represent a selected cell population having a higher average affinity for antigen than the starting pool of antigen-responsive naïve cells.

As discussed above, memory T cells are phenotypically heterogeneous. While the extent to which this heterogeneity also applies to the functional properties of memory cells remains to be fully investigated, it is evident that central memory cells and effector memory cells exhibit clear differences in their response to antigen. Thus, a direct comparison of these two subpopulations showed that CD4<sup>+</sup> CD45R0<sup>+</sup> CCR7<sup>+</sup> effector memory cells secrete a broad range of cytokines (IL-2, IFN- $\gamma$ , IL-4 and IL-5) within 24 hr of stimulation through the TCR, while CD4<sup>+</sup> CD45R0<sup>+</sup> CCR7<sup>+</sup> central memory cells secrete only IL-2.<sup>17</sup> In addition, effector memory cells were also able to proliferate in response to lower concentrations of anti-CD3 antibodies than central memory cells, although the latter were still more responsive than naïve T cells. Therefore, memory T cells are in fact functionally heterogeneous, and any general distinction between the functional properties of naïve and memory T cells is clearly an oversimplification. This heterogeneity may partially account for discrepancies in the literature regarding differences between naïve and memory cells.

## MIGRATION OF NAÏVE AND MEMORY T CELLS

Lymphocyte migration is controlled by the combination of adhesion molecules and chemokine receptors expressed by lymphocytes. Since the expression of these molecules differs between naïve and memory T cells, it is not surprising that these cells have different migratory properties.

Naïve T cells exhibit a restricted pattern of migration, in which they move continuously between the secondary lymphoid organs (spleen, lymph nodes and Peyer's patches) via blood and lymph; transit through individual organs takes about 12–18 hr.<sup>111</sup> The expression of CD62L and CCR7 by naïve T cells plays a key role in establishing this recirculation pattern. Thus, entry of naïve T cells into lymph nodes and Peyer's patches occurs at HEVs in a CD62L- and CCR7-dependent manner (see above). In addition, although initial entry of lymphocytes into the spleen is a passive process, with T cells being deposited from the blood into the marginal zone, migration of cells into the T-cell zones of the spleen may also require expression of CCR7.<sup>112</sup> However, naïve T cells lack homing receptors for peripheral tissues and chemokine receptors for inflammatory cytokines and are therefore unable to enter non-lymphoid tissues.

By contrast, some memory T cells express chemokine and adhesion receptors that enable them to extravasate into non-lymphoid tissues.<sup>17</sup> Migration into peripheral tissues was originally demonstrated for memory-phenotype T cells in sheep and subsequently demonstrated for antigen-specific memory T cells in mice.<sup>6,59,113–116</sup> However, as discussed above, memory T cells are heterogeneous with respect to the expression of homing molecules and likewise exhibit heterogeneity in migration. Central memory cells, which express CD62L and CCR7, are able to enter lymph nodes via HEVs and exhibit a pattern of migration that is similar to that of naïve T cells. Conversely, effector memory cells lack CD62L and CCR7 but express homing receptors that allow them to enter non-lymphoid tissues; these cells may reach lymph nodes via afferent lymph rather than through HEVs.<sup>6</sup> In addition, subpopulations of memory T cells exhibit a preference for migration to particular tissues, for example gut vs. skin, on the basis of the particular homing molecules expressed.<sup>113,117–122</sup> This predilection for specific tissues may be dictated by the local environment in which initial T-cell priming occurs.<sup>121–123</sup>

The migration pathways favoured by naïve and memory T cells are linked to their activation requirements and functional properties. As described above, activation of naïve T cells requires antigen presentation on mature DCs. These APCs are found in the T-cell zones of secondary lymphoid organs; DCs move into these areas and present antigen following the uptake of pathogens in peripheral sites.<sup>124</sup> Hence, secondary lymphoid organs serve as predetermined meeting places for naïve T cells and DCs that maximize the chance of low-frequency T cells encountering their specific antigen. Continuous movement of naïve T cells between different lymphoid organs allows these cells to scan the surface of DCs derived from all parts of the body. In addition, the environment of the organised lymphoid tissues

is ideal for expansion of activated T cells and their differentiation into effectors.

Conversely, activation of memory T cells can occur outside the optimized conditions of the secondary lymphoid organs, owing both to the increased frequency of antigen-specific cells among memory cells and to their ability to become activated by a wider range of APCs. As a consequence, effector memory T cells can efficiently perform a surveillance function at potential sites of pathogen invasion and provide a rapid response to re-infection. In contrast, central memory T cells, which appear to migrate in a similar manner to naïve T cells, probably respond to antigen in secondary lymphoid organs. Whether central memory cells actually require antigen presentation on DCs in order to undergo activation and differentiation into effector cells is unclear (see above). Nevertheless, it seems likely that a much greater degree of expansion will occur following antigenic stimulation in lymphoid organs. Therefore, central and effector memory T cells may provide complementary functions upon secondary infection, with effector memory cells providing an immediate local response and central memory cells rapidly generating large numbers of effectors.

## LIFESPAN OF NAÏVE VS. MEMORY T CELLS

In general, cell populations can be maintained at constant numbers by two mechanisms: (1) survival of long-lived cells that do not divide, and (2) proliferation of shorter lived cells, which is balanced by the same rate of cell death (turnover). Under non-disease conditions, the overall rate of T-cell turnover is relatively slow, indicating that these cells can remain in interphase for long periods of time. This was shown initially in rodents, where T-cell proliferation *in vivo* was measured by infusion of a labelled DNA precursor, <sup>3</sup>H-thymidine.<sup>125–129</sup> From this type of study, the turnover time for T cells in the thoracic duct of mice was calculated to be of the order of 4–6 months.<sup>127</sup> More recent studies using <sup>2</sup>H-glucose as a DNA precursor have shown that human T cells also turn over slowly.<sup>130</sup> Here, the mean intermitotic times for CD4<sup>+</sup> and CD8<sup>+</sup> T cells in peripheral blood were estimated to be 87 and 77 days, respectively.

Studies in a number of different species, including mice,<sup>131</sup> sheep,<sup>6</sup> rhesus macaques<sup>132</sup> and humans,<sup>133,134</sup> have shown that memory-phenotype T cells turn over at a faster rate than naïve-phenotype T cells. On average, the rate of incorporation of labelled DNA precursors is approximately 5 times faster for memory-phenotype cells than naïve-phenotype cells. Extrapolation of these data directly to naïve and memory T cells is complicated by the limitations of using phenotypic markers to identify these cells (see above). However, it is notable that studies of cell populations that can be designated as memory T cells with high confidence – cells analysed at extended periods after immunization, using TCR transgenics or MHC tetramers to identify antigen-specific cells – have shown that these cells exhibit similar rates of division *in vivo* to memory-phenotype T cells in normal mice.<sup>55,135,136</sup> The implication therefore is that memory T cells are maintained by



intermittent cell division while naïve T cells persist as relatively quiescent, non-dividing cells.

Although memory-phenotype T cells exhibit rapid turnover overall, some memory T cells appear to remain in a non-dividing state for long periods of time. This is evident from the results of DNA labelling studies, which show considerable heterogeneity in the rates at which cells incorporate label during the infusion phase or lose label after the termination of treatment.<sup>131,134</sup> This is not surprising in view of the evidence cited above for heterogeneity amongst memory T cells. At present, it is unknown whether long- and short-lived memory T cells correspond to subpopulations of cells that can be distinguished phenotypically (e.g. central vs. effector memory T cells).

#### FACTORS CONTROLLING THE MAINTENANCE OF NAÏVE AND MEMORY T CELLS

In addition to exhibiting distinct kinetic behaviour, naïve and memory T cells are also dependent on different factors for maintaining their survival. Two main stimuli have been studied in this regard: (1) MHC-peptide, and (2) cytokines.

##### MHC-peptide

The issue of whether long-term survival of naïve T cells is dependent on contact with self-MHC-peptide ligands has been studied extensively.<sup>55,137–145</sup> Because MHC expression in the thymus is essential for the initial development of T cells, investigating the role of peripheral MHC in T-cell survival requires either adoptive transfer of mature T cells into MHC-deficient hosts or selective and transient expression of MHC in the thymus of such mice. In general, these studies have shown that the lifespan of naïve CD4<sup>+</sup> and CD8<sup>+</sup> T cells is shortened in the absence of MHC class II or class I, respectively. However, it is notable that naïve CD4<sup>+</sup> T cells can still survive for considerable periods of time in the absence of MHC class II. Thus, the half-life of naïve CD4<sup>+</sup> cells in the absence of MHC class II has been estimated to be approximately 3–4 weeks by some investigators,<sup>140,145</sup> while others have reported that naïve CD4 cells can survive equally well in control and MHC class II-deficient hosts for the first 4<sup>144</sup> or 8<sup>143</sup> weeks after transfer. These results imply that naïve CD4 cells require only very infrequent contact with MHC in order to survive.

The nature of the MHC-ligands involved in transmitting survival signals to naïve T cells is unknown. For homeostatic proliferation (see above), T cells respond to self-MHC in a peptide-specific manner, with the peptides involved being at least partially related to those involved in T-cell positive selection in the thymus.<sup>144,146–148</sup> Therefore, one possibility is that the same MHC-peptide ligands that stimulate proliferation in lymphopenic animals trigger survival under normal conditions. Conversely, survival of naïve T cells may depend only upon recognition of MHC and be independent of the bound peptide.

There has been extensive debate surrounding the role of persisting antigen in the maintenance of T-cell memory. While it remains a contentious issue whether periodic

contact with antigen is required to maintain protective immunity,<sup>149–153</sup> it is now generally accepted that the survival of memory T cells is antigen-independent. Thus, numerous studies in mice have shown that both CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells of defined specificity survive indefinitely following adoptive transfer into antigen-free hosts.<sup>54,55,91–93,154–157</sup> Furthermore, long-term survival also applies to CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells transferred to MHC class II- and MHC class I-deficient mice, respectively.<sup>55,157–159</sup> Therefore, unlike naïve T cells, memory T cells do not appear to depend on interactions with MHC for survival. Notably, memory T cells also continue to proliferate in MHC-deficient hosts.<sup>55,141,157,159</sup> For CD8<sup>+</sup> memory T cells, proliferation occurs at similar rates after transfer to MHC class I<sup>+/+</sup> or MHC class I<sup>-/-</sup> hosts, implying that turnover, like survival is regulated by signals independent of TCR triggering.<sup>55</sup>

By contrast, it is less clear whether the functional properties of memory T cells are also independent of contact with MHC. For some properties of memory T cells, this does appear to be the case. Thus, CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells retain a CD44<sup>hi</sup> phenotype and the capacity for rapid secretion of IFN- $\gamma$  after transfer to MHC-deficient mice.<sup>55,159</sup> However, it has been reported that CD4 memory cells parked in MHC class II-deficient hosts lose other functional characteristics ascribed to memory cells, namely the ability to respond to antigen presented by non-professional APCs such as B cells and a relative independence from costimulation.<sup>159</sup> Whether a similar requirement for contact with MHC exists for CD8 memory cells remains to be investigated.

##### Cytokines

The main cytokine that has been implicated in maintaining the survival of naïve T cells is IL-7. In fact, IL-7 seems to be essential for naïve T cell survival, as these cells disappear rapidly following adoptive transfer to IL-7<sup>-/-</sup> recipients or in normal mice treated with anti-IL-7R antibodies; in addition, IL-7R<sup>-/-</sup> naïve T cells fail to survive in normal hosts.<sup>70,160,161</sup> Although other cytokines such as IL-4, IL-6 and IL-15 have been shown to rescue naïve T cells from apoptosis *in vitro*<sup>162,163</sup> (M. Berard and D. F. Tough, unpublished data), these cytokines do not appear to be required for the survival of naïve T cells *in vivo* and cannot compensate for the absence of IL-7.<sup>161</sup> In addition to promoting the survival of naïve T cells in normal mice, IL-7 also seems to be required for homeostatic proliferation of naïve T cells under lymphopenic conditions.<sup>70,160</sup> Thus, maintenance of naïve T cells appears to be strictly dependent on contact with both self-MHC-peptide ligands and IL-7.

Evidence that cytokines play a role in the maintenance of memory T cells came initially from studies showing that injection of either type I IFN (IFN- $\alpha/\beta$ ) or inducers of IFN- $\alpha/\beta$  stimulated TCR-independent proliferation of memory-phenotype (CD44<sup>hi</sup>) CD8<sup>+</sup> T cells *in vivo*.<sup>164,165</sup> Subsequently, a number of other cytokines, including IL-12, IL-15, IL-18 and IFN- $\gamma$ , were found to have similar

effects.<sup>18,166</sup> Notably, enhanced proliferation was restricted to CD44<sup>hi</sup> CD8<sup>+</sup> T cells following injection of each of these cytokines; little or no cell division was induced amongst naïve phenotype T cells. Of the cytokines shown to promote proliferation of memory-phenotype CD8<sup>+</sup> T cells *in vivo*, however, only IL-15 was able to stimulate these cells to divide when added to purified T cells *in vitro*. Here, the selective proliferation of memory-phenotype CD8<sup>+</sup> T cells in response to IL-15 was associated with much higher expression of the IL-15R  $\beta$ -chain (CD122) on CD44<sup>hi</sup> vs. CD44<sup>lo</sup> CD8<sup>+</sup> T cells.<sup>18</sup> Since IFN- $\alpha/\beta$ , IFN- $\gamma$ , IL-12 and IL-18 are all capable of up-regulating IL-15 expression by APCs (the latter two through induction of IFN- $\gamma$ ), it has been proposed that IL-15 acts as a final common effector molecule mediating the effects on CD44<sup>hi</sup> CD8<sup>+</sup> T cell proliferation.<sup>18,166</sup>

Based on the observation that injection of IL-15 or induction of IL-15 expression *in vivo* results in enhanced proliferation of memory-phenotype CD8<sup>+</sup> T cells, it was hypothesized that the high proliferation of these cells in normal mice might be attributable to background production of IL-15. In support of this idea, injection of anti-CD122 antibodies into mice was shown to strongly reduce the proliferation of CD44<sup>hi</sup> CD8<sup>+</sup> T cells.<sup>167</sup> Although this antibody can block signalling through both the IL-15R and the IL-2R, the inhibitory effects appeared to result from blocking IL-15, since injection of anti-IL-2 + anti-IL-2R $\alpha$  actually enhanced CD44<sup>hi</sup> CD8<sup>+</sup> T-cell proliferation. The conclusion from this study was that IL-15 and IL-2 had opposing effects on the proliferation of memory-phenotype CD8 T cells *in vivo*, with IL-15 enhancing and IL-2 inhibiting cell division.

Strikingly, the number of memory-phenotype CD8<sup>+</sup> T cells is markedly reduced in both IL-15- and IL-15R $\alpha$ -deficient mice,<sup>168,169</sup> and increased in transgenic mice over-expressing IL-15.<sup>170,171</sup> Thus, IL-15 appears to play an important role in regulating the number of CD44<sup>hi</sup> CD8<sup>+</sup> T cells found in normal mice; definitive evidence of its importance for antigen-specific CD8 memory T cells awaits further study in infection/immunization models. Exactly how IL-15 controls the maintenance of CD44<sup>hi</sup> CD8<sup>+</sup> T cells remains unclear. The simplest idea is that IL-15-induced proliferation is necessary to counter a rapid rate of death; increases or decreases in the amount of IL-15 tip the balance towards more or less accumulation of CD44<sup>hi</sup> CD8<sup>+</sup> T cells, respectively. However, the fact that a considerable proportion of memory-phenotype CD8<sup>+</sup> T cells can persist for long periods of time in a non-dividing state is inconsistent with the idea that all memory cells are intrinsically short-lived.<sup>131</sup> Therefore, an alternative possibility is that IL-15 is able to promote survival as well as proliferation of CD44<sup>hi</sup> CD8<sup>+</sup> T cells. In this respect, it is notable that IL-15 induces up-regulated expression of the pro-survival factor Bcl-2 in CD8<sup>+</sup> T cells.<sup>171</sup>

In contrast to its effects on CD44<sup>hi</sup> CD8<sup>+</sup> T cells, IL-15 stimulates little proliferation of memory-phenotype CD4<sup>+</sup> T cells either *in vivo* or *in vitro*.<sup>18</sup> Furthermore, memory-phenotype CD4<sup>+</sup> T cells are found in normal numbers in IL-15<sup>-/-</sup> and IL-15R $\alpha$ <sup>-/-</sup> mice.<sup>168,169</sup> Therefore, IL-15


does not appear to play a role in the maintenance of CD4<sup>+</sup> memory cells. In fact, CD4<sup>+</sup> memory cells are long-lived in mice deficient for the common  $\gamma$ -chain, indicating that none of the cytokines utilizing this receptor component (IL-2, IL-4, IL-7, IL-9 and IL-15) are essential for the survival of CD4<sup>+</sup> memory T cells.<sup>71</sup> Thus, if cytokines do play a role in the maintenance of CD4<sup>+</sup> memory T cells, the factors involved are clearly distinct from those that have effects on either naïve T cells or CD8<sup>+</sup> memory cells. Presently, the best evidence that the turnover of CD4<sup>+</sup> memory T cells can be regulated by cytokines comes from a study in which T-cell proliferation was assessed following the activation of natural killer (NK1.1<sup>+</sup>) T cells *in vivo*.<sup>172</sup> Under these conditions, the proliferation of CD44<sup>hi</sup> CD4<sup>+</sup> (NK1.1<sup>-</sup>) T cells was markedly increased through a mechanism that was dependent on IL-12 or IFN- $\gamma$ .

## CONCLUSIONS

There is considerable evidence to support the idea that T cells retain a permanent imprint of a prior response to antigen. In broad terms, naïve and memory T cells differ in phenotype, pattern of migration, responsiveness to antigen and cytokines and requirements for survival. However, it is also clear that memory T cells are heterogeneous with respect to all of these parameters. The recent identification of markers that can be used to separate subpopulations of memory T cells will allow a more precise delineation of the characteristics of memory cells.

Variation amongst memory T cells may arise as a result of the initial conditions of activation, but might also be influenced by the environment in which memory T cells reside. In this respect, future work aimed at elucidating the relationship between phenotypically distinct subpopulations of memory T cells should provide important insight into the nature of T-cell memory. For example, while it is evident that CD4<sup>+</sup> central memory cells can differentiate into cells resembling effector memory cells upon stimulation with antigen or cytokines *in vitro*, the rate at which this conversion takes place *in vivo*, and whether reversion from effector to central memory cell will occur in the absence of such stimulation, are not known.<sup>17,173</sup> In addition, the identification of memory T cells expressing certain markers that are characteristic of naïve T cells raises the question of whether some memory T cells may in fact be completely indistinguishable from naïve T cells in phenotype. If so, it would be of interest to determine whether these cells retain any functional characteristics of memory cells.

Finally, the extent to which the altered functional properties of memory T cells contribute to the characteristics of the secondary immune response remains unclear. In practice, changes at the single cell level probably synergize with increases in frequency in producing a rapid response to secondary infection (Fig. 1). Hence, effector memory T cells are able to respond immediately to re-infection on the basis of their distribution, reduced activation requirements and increased frequency, and should therefore limit the early spread of infection.<sup>116</sup>



**Figure 1.** Combined effects of qualitative and quantitative changes in memory T cells in mediating a rapid secondary response to infection. Effector memory T cells are situated in peripheral tissues at sites of pathogen entry. These cells exhibit effector function immediately upon recognition of antigen presented on non-professional APCs and limit the early spread of infection. Central memory T cells are activated in secondary lymphoid organs (i.e. the draining lymph node) following recognition of antigen on DCs and generate large numbers of effector cells. The response of central memory cells is prompt as a result of both a high frequency of antigen specific cells and their ability to differentiate rapidly into effectors.

At the same time, central memory T cells are capable of generating rapidly a large number of effector cells based on their high precursor frequency; an accelerated ability to differentiate into effectors may also play a role.<sup>174</sup> These effector cells then distribute systemically and act to clear the residual infection. Understanding how memory T cells are maintained not only at a high frequency but also in an enhanced state of functional readiness therefore is important for both our fundamental understanding of the immune system and for the rational design of vaccines.

#### ACKNOWLEDGMENTS

This is publication number 56 from the Edward Jenner Institute for Vaccine Research.

#### REFERENCES

- 1 Budd RC, Cerottini J-C, MacDonald HR. Phenotypic identification of memory cytolytic T lymphocytes in a subset of Ly-2+ cells. *J Immunol* 1987; **138**:1009–13.
- 2 Budd RC, Cerottini JC, MacDonald HR. Selectively increased production of interferon-gamma by subsets of Lyt2+ and L3T4+ T cells identified by expression of Pgp-1. *J Immunol* 1987; **138**:3583–6.
- 3 Tabi Z, Lynch F, Ceredig R, Allan JE, Doherty PC. Virus-specific memory T cells are Pgp-1+ and can be selectively activated with phorbol ester and calcium ionophore. *Cell Immunol* 1988; **113**:268–77.
- 4 Sanders ME, Makgoba MW, Sharrow SO, Stephany D, Springer TA, Young HA, Shaw S. Human memory T lymphocytes express increased levels of three cell adhesion

- molecules (LFA-3, CD2, and LFA-1) and three other molecules (UCHL1, CDw29, and Pgp-1) and have enhanced IFN- $\gamma$  production. *J Immunol* 1988; **140**:1401–7.
- 5 Butterfield K, Fathman CG, Budd RC. A subset of memory CD4+ helper T lymphocytes identified by expression of Pgp-1. *J Exp Med* 1989; **169**:1461–6.
- 6 Mackay CR, Marston WL, Dudler L. Naive and memory T cells show distinct pathways of lymphocyte recirculation. *J Exp Med* 1990; **171**:801–17.
- 7 Buckle AM, Hogg N. Human memory T cells express intercellular adhesion molecule-1 which can be increased by interleukin 2 and interferon- $\gamma$ . *Eur J Immunol* 1990; **20**:337–41.
- 8 Beverley PCL. Functional analysis of human T cell subsets defined by CD45 isoform expression. *Sem Immunol* 1992; **4**:35–41.
- 9 McFarland HI, Nahill SR, Maciaszek JW, Welsh RM. CD11b (Mac-1): a marker for CD8+ cytotoxic T cell activation and memory in virus infection. *J Immunol* 1992; **149**:1326–33.
- 10 Okumura M, Fujii Y, Takeuchi Y, Inada K, Nakahara K, Matsuda H. Age-related accumulation of LFA-1<sup>high</sup> cells in a CD8+ CD45RA<sup>high</sup> T cell population. *Eur J Immunol* 1993; **23**:1057–63.
- 11 Hamann D, Baars PA, Rep MHG, Hooibrink B, Kerkhof-Garde SR, Klein MR, van Lier RAW. Phenotypic and functional separation of memory and effector human CD8+ T cells. *J Exp Med* 1997; **186**:1407–18.
- 12 Berg EL, Robinson MK, Warnock RA, Butcher EC. The human peripheral lymph node vascular addressin is a ligand for LECAM-1, the peripheral lymph node homing receptor. *J Cell Biol* 1991; **114**:343–9.
- 13 Gunn MD, Tangemann K, Tam C, Cyster JG, Rosen SD, Williams LT. A chemokine expressed in lymphoid high endothelial venules promotes the adhesion and chemotaxis of naive T lymphocytes. *Proc Natl Acad Sci USA* 1998; **95**:258–63.
- 14 Swain SL, Bradley LM, Croft M *et al.* Helper T-cell subsets: phenotype, function and the role of lymphokines in regulating their development. *Immunol Rev* 1991; **123**:115–44.
- 15 Bradley LM, Atkins GG, Swain SL. Long-term CD4+ memory T cells from the spleen lack MEL-14, the lymph node homing receptor. *J Immunol* 1992; **148**:324–31.
- 16 Hou S, Doherty PC. Partitioning of responder CD8+ T cells in lymph node and lung of mice with sendai virus pneumonia by LECAM-1 and CD45RB phenotype. *J Immunol* 1993; **150**:5494–500.
- 17 Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 1999; **401**:708–12.
- 18 Zhang X, Sun S, Hwang I, Tough DF, Sprent J. Potent and selective stimulation of memory-phenotype CD8+ T cells in vivo by IL-15. *Immunity* 1998; **8**:591–9.
- 19 Cho BK, Wang C, Sugawa S, Eisen HN, Chen J. Functional differences between memory and naive CD8 T cells. *Proc Natl Acad Sci USA* 1999; **96**:2976–81.
- 20 Walunas TL, Bruce DS, Dustin L, Loh DY, Bluestone JA. Ly-6C is a marker of memory CD8+ T cells. *J Immunol* 1995; **155**:1873–83.
- 21 Curtsinger JM, Lins DC, Mescher MF. CD8+ memory T cells (CD44<sup>high</sup>, Ly-6C<sup>+</sup>) are more sensitive than naive cells (CD44<sup>low</sup>, Ly-6C<sup>-</sup>) to TCR/CD8 signaling in response to antigen. *J Immunol* 1998; **160**:3236–43.
- 22 Merckenschlager M, Terry L, Edwards R, Beverley PCL. Limiting dilution analysis of proliferative responses in human



- lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. *Eur J Immunol* 1988; **18**:1653–61.
- 23 Merkenschlager M, Beverley PCL. Evidence for differential expression of CD45 isoforms by precursors for memory-dependent and independent cytotoxic responses: human CD8 memory CTLp selectively express CD45RO (UCHL1). *Int Immunol* 1989; **1**:450–9.
  - 24 Hayward AR, Lee J, Beverley PC. Ontogeny of expression of UCHL1 antigen on TcR + (CD4/8) and TCR  $\delta$  + T cells. *Eur J Immunol* 1989; **19**:189.
  - 25 Birkeland ML, Johnson P, Trowbridge IS, Pure E. Changes in CD45 isoform expression accompany antigen-induced murine T-cell activation. *Proc Natl Acad Sci USA* 1989; **86**:6734–8.
  - 26 Powrie F, Mason D. The MRC OX-22- CD4+ T cells that help B cells in secondary immune responses derive from naive precursors with the MRC OX-22+ CD4+ phenotype. *J Exp Med* 1989; **169**:653–62.
  - 27 Lee WT, Yin XM, Vitetta ES. Functional and ontogenic analysis of murine CD45R<sup>hi</sup> and CD45R<sup>lo</sup> CD4+ T-cells. *J Immunol* 1990; **144**:3288–95.
  - 28 Marvel J, Lightstone E, Samberg NL, Ettinghausen D, Stauss HJ. The CD45RA molecule is expressed in naive murine CTL precursors but absent in memory and effector CTL. *Int Immunol* 1991; **3**:21–8.
  - 29 Waldmann TA, Tagaya Y. The multifaceted regulation of interleukin-15 expression and the role of this cytokine in NK cell differentiation and host response to intracellular pathogens. *Annu Rev Immunol* 1999; **17**:19–49.
  - 30 Gumley TP, McKenzie IF, Sandrin MS. Tissue expression, structure and function of the murine Ly-6 family of molecules. *Immunol Cell Biol* 1995; **73**:277–96.
  - 31 Trowbridge IS, Thomas ML. CD45: an emerging role as a protein tyrosine phosphatase required for lymphocyte activation and development. *Annu Rev Immunol* 1994; **12**:85–116.
  - 32 Irie-Sasaki J, Sasaki T, Matsumoto W *et al.* CD45 is a JAK phosphatase and negatively regulates cytokine receptor signalling. *Nature* 2001; **409**:349–54.
  - 33 Chervenak R, Cohen JJ. Peanut lectin binding as a marker for activated T-lineage lymphocytes. *Thymus* 1982; **4**:61–7.
  - 34 Piller F, Piller V, Fox RI, Fukuda M. Human T-lymphocyte activation is associated with changes in O-glycan biosynthesis. *J Biol Chem* 1988; **263**:15146–50.
  - 35 Galvan M, Murali-Krishna K, Ming LL, Baum L, Ahmed R. Alterations in cell surface carbohydrates on T cells from virally infected mice can distinguish effector/memory CD8+ T cells from naive cells. *J Immunol* 1998; **161**:641–8.
  - 36 Priatel JJ, Chui D, Hiraoka N *et al.* The ST3Gal-I sialyltransferase controls CD8+ T lymphocyte homeostasis by modulating O-glycan biosynthesis. *Immunity* 2000; **12**:273–83.
  - 37 Harrington LE, Galvan M, Baum LG, Altman JD, Ahmed R. Differentiating between memory and effector CD8 T cells by altered expression of cell surface O-glycans. *J Exp Med* 2000; **191**:1241–6.
  - 38 Stout RD, Suttles J. T cells bearing the CD44<sup>hi</sup> 'memory' phenotype display characteristics of activated cells in G1 stage of cell cycle. *Cell Immunol* 1992; **141**:433–43.
  - 39 Bell EB, Sparshott SM. Interconversion of CD45R subsets of CD4 T cells *in vivo*. *Nature* 1990; **348**:163–6.
  - 40 Gahring LC, Ernst DN, Romball CG, Thoman ML, Torbett BE, Hobbs M, Weigle WO. The expression of CD45RB on antigen-responsive CD4+ lymphocytes: mouse strain polymorphism and different responses to distinct antigens. *Cell Immunol* 1993; **148**:269–82.
  - 41 Sparshott SM, Bell EB. Membrane CD45R isoform exchange on CD4 T cells is rapid, frequent and dynamic *in vivo*. *Eur J Immunol* 1994; **24**:2573–8.
  - 42 Pihlgren M, Lightstone L, Mamalaki C, Rimón G, Kioussis D, Marvel J. Expression *in vivo* of CD45RA, CD45RB and CD44 on T cell receptor-transgenic CD8+ T cells following immunization. *Eur J Immunol* 1995; **25**:1755–9.
  - 43 Pilling D, Akbar AN, Bacon PA, Salmon M. CD4+ CD45RA+ T cells from adults respond to recall antigens after CD28 ligation. *Int Immunol* 1996; **8**:1737–42.
  - 44 Richards D, Chapman MD, Sasama J, Lee TH, Kemeny DM. Immune memory in CD4+ CD45RA+ T cells. *Immunology* 1997; **91**:331–9.
  - 45 Wilamasundera S, Katz DR, Chain BM. Responses to human rhinovirus in CD45 T cell subsets isolated from tonsil. *Eur J Immunol* 1998; **28**:4374–81.
  - 46 Faint JM, Annels NE, Curnow SJ *et al.* Memory T cells constitute a subset of the human CD8+CD45RA+ pool with distinct phenotypic and migratory characteristics. *J Immunol* 2001; **167**:212–20.
  - 47 Jung TM, Gallatin WM, Weissman IL, Dailey MO. Downregulation of homing receptors after T cell activation. *J Immunol* 1988; **141**:4110–7.
  - 48 Rothstein DM, Yamada A, Schlossman SF, Morimoto C. Cyclic regulation of CD45 isoform expression in a long term human CD4+CD45RA+ T cell line. *J Immunol* 1991; **146**:1175–83.
  - 49 Warren HS, Skipsey LJ. Loss of activation-induced CD45RO with maintenance of CD45RA expression during prolonged culture of T cells and NK cells. *Immunology* 1991; **74**:78–85.
  - 50 Fujii Y, Okumura M, Inada K, Nakahara K. Reversal of CD45R isoform switching in CD8+ T cells. *Cell Immunol* 1992; **139**:176–84.
  - 51 Tripp RA, Hou S, Doherty PC. Temporal loss of the activated 1-selectin-low phenotype for virus-specific CD8+ memory T cells. *J Immunol* 1995; **154**:5870–5.
  - 52 Andersen P, Smedegaard B. CD4+ T-cell subsets that mediate immunological memory to *Mycobacterium tuberculosis* infection in mice. *Infect Immun* 2000; **68**:621–9.
  - 53 Boursalian RE, Bottomly K. Stability of naive and memory phenotypes on resting CD4 T cells *in vivo*. *J Immunol* 1999; **162**:9–16.
  - 54 Lau LL, Jamieson BD, Somasundaram R, Ahmed R. Cytotoxic T cell memory without antigen. *Nature* 1994; **369**:648–52.
  - 55 Murali-Krishna K, Lau LL, Sambhara S, Lemonnier F, Altman J, Ahmed R. Persistence of memory CD8 T cells in MHC class I-deficient mice. *Science* 1999; **286**:1377–81.
  - 56 Nociari MM, Telford W, Russo C. Postthymic development of CD28-CD8+ T cell subset: age-associated expansion and shift from memory to naive phenotype. *J Immunol* 1999; **162**:3327–35.
  - 57 Wikby A, Johansson B, Olsson J, Lofgren S, Nilsson BO, Ferguson F. Expansions of peripheral blood CD8 T-lymphocyte subpopulations and an association with cytomegalovirus seropositivity in the elderly: the Swedish NONA immune study. *Exp Gerontol* 2002; **37**:445–53.
  - 58 Manjunath N, Shankar P, Wan J *et al.* Effector differentiation is not prerequisite for generation of memory cytotoxic T lymphocytes. *J Clin Invest* 2001; **108**:871–8.



- 59 Weninger W, Crowley MA, Manjunath N, von Andrian UH. Migratory properties of naïve, effector, and memory CD8<sup>+</sup> T cells. *J Exp Med* 2001; **194**:953–66.
- 60 Goldrath AW, Bevan MJ. Low-affinity ligands for the TCR drive proliferation of mature CD8<sup>+</sup> T cells in lymphopenic hosts. *Immunity* 1999; **11**:183–90.
- 61 Ernst B, Lee D-S, Chang JM, Sprent J, Surh CD. The peptide ligands mediating positive selection in the thymus control T cell survival and homeostatic proliferation in the periphery. *Immunity* 1999; **11**:173–81.
- 62 Kieper WC, Jameson SC. Homeostatic expansion and phenotypic conversion of naïve T cells in response to self peptide/MHC ligands. *Proc Natl Acad Sci USA* 1999; **96**:13306–11.
- 63 Oehen S, Brduscha-Riem K. Naïve cytotoxic T lymphocytes spontaneously acquire effector function in lymphocytopenic recipients: a pitfall for T cell memory studies? *Eur J Immunol* 1999; **29**:608–14.
- 64 Viret C, Wong FS, Janeway CA Jr. Designing and maintaining the mature TCR repertoire: the continuum of self-peptide: self-MHC complex recognition. *Immunity* 1999; **10**:559–68.
- 65 Goldrath AW, Bogatzki LY, Bevan MJ. Naïve T cells transiently acquire a memory-like phenotype during homeostasis-driven proliferation. *J Exp Med* 2000; **192**:557–64.
- 66 Cho BK, Rao VP, Ge Q, Eisen HN, Chen J. Homeostasis-stimulated proliferation drives naïve T cells to differentiate directly into memory T cells. *J Exp Med* 2000; **192**:549–56.
- 67 Murali-Krishna K, Ahmed R. Naïve T cells masquerading as memory cells. *J Immunol* 2000; **165**:1733–7.
- 68 Muranski P, Chmielowski B, Ignatowicz L. Mature CD4<sup>+</sup> T cells perceive a positively selecting class II MHC/peptide complex in the periphery. *J Immunol* 2000; **164**:3087–94.
- 69 Boursalian TE, Bottomly K. Survival of naïve CD4 T cells: roles of restricting versus selecting MHC class II and cytokine milieu. *J Immunol* 1999; **162**:3795–801.
- 70 Schluns KS, Kieper WC, Jameson SC, Lefrançois L. Interleukin-7 mediates the homeostasis of naïve and memory CD8 T cells *in vivo*. *Nature Immunol* 2000; **1**:426–32.
- 71 Lantz O, Grandjean I, Matzinger P, Di Santo JP.  $\gamma$  chain required for naïve CD4<sup>+</sup> T cell survival but not for antigen proliferation. *Nature Immunol* 2000; **1**:54–8.
- 72 Engers HD, MacDonald HR. Generation of cytolytic T lymphocytes *in vitro*. *Contemp Top Immunobiol* 1976; **5**:145–90.
- 73 Byrne JA, Butler JL, Cooper MD. Differential activation requirements for virgin and memory T cells. *J Immunol* 1988; **141**:3249–57.
- 74 Byrne JA, Butler JL, Reinherz EL, Cooper MD. Virgin and memory T cells have different requirements for activation via the CD2 molecule. *Int Immunol* 1989; **1**:29–35.
- 75 Sanders ME, Makgoba MW, June CH, Young HA, Shaw S. Enhanced responsiveness of human memory T cells to CD2 and CD3 receptor-mediated activation. *Eur J Immunol* 1989; **19**:803–8.
- 76 Ehlers S, Smith KA. Differentiation of T cell lymphokine gene expression: the *in vitro* acquisition of T cell memory. *J Exp Med* 1991; **173**:25–36.
- 77 Akbar AN, Salmon M, Janosy G. The synergy between naïve and memory T cells during activation. *Immunol Today* 1991; **12**:184–8.
- 78 de Jong R, Brouwer M, Miedema F, van Lier RA. Human CD8<sup>+</sup> T lymphocytes can be divided into CD45RA<sup>+</sup> and CD45RO<sup>+</sup> cells with different requirements for activation and differentiation. *J Immunol* 1991; **146**:2088–94.
- 79 Luqman M, Bottomly K. Activation requirements for CD4<sup>+</sup> T cells differing in CD45R expression. *J Immunol* 1992; **149**:2300–6.
- 80 Fischer H, Gyorloff A, Hedlund G, Hedman H, Lundgren E, Kalland T, Sjogren HO, Dohlsten M. Stimulation of human naïve and memory T helper cells with bacterial superantigen. Naïve CD4<sup>+</sup>45RA<sup>+</sup> T cells require a costimulatory signal mediated through the LFA-1/ICAM-1 pathway. *J Immunol* 1992; **148**:1993–8.
- 81 Kuiper H, Brouwer M, de Boer M, Parren P, van Lier RAW. Differences in responsiveness to CD3 stimulation between naïve and memory CD4<sup>+</sup> T cells cannot be overcome by CD28 costimulation. *Eur J Immunol* 1994; **24**:1956–60.
- 82 Kearney ER, Pape KA, Loh DY, Jenkins MK. Visualization of peptide-specific T cell immunity and peripheral tolerance induction *in vivo*. *Immunity* 1994; **1**:327–39.
- 83 Pihlgren M, Dubois PM, Tomkowiak M, Sjogren T, Marvel J. Resting memory CD8<sup>+</sup> T cells are hyperreactive to antigenic challenge *in vitro*. *J Exp Med* 1996; **184**:2141–51.
- 84 Rogers PR, Dubey C, Swain SL. Qualitative changes accompany memory T cell generation. Faster, more effective responses at lower doses of antigen. *J Immunol* 2000; **164**:2338–46.
- 85 Slifka MK, Whitton JL. Functional avidity maturation of CD8<sup>+</sup> T cells without selection of higher affinity TCR. *Nature Immunol* 2001; **2**:711–7.
- 86 Constant S, Zain M, West J, Pasqualini T, Ranney P, Bottomly K. Are primed CD4<sup>+</sup> T lymphocytes different from unprimed cells? *Eur J Immunol* 1994; **24**:1073–9.
- 87 Bachmann MF, Barner M, Viola A, Kopf M. Distinct kinetics of cytokine production and cytotoxicity in effector and memory T cells after viral infection. *Eur J Immunol* 1999; **29**:291–9.
- 88 Zimmermann C, Prevost-Blondel A, Blaser C, Pircher H. Kinetics of the response of naïve and memory CD8 T cells to antigen: similarities and differences. *Eur J Immunol* 1999; **29**:284–90.
- 89 Inaba K, Metlay JP, Crowley MT, Witmer-Pack M, Steinman RM. Dendritic cells as antigen presenting cells *in vivo*. *Int Rev Immunol* 1990; **6**:197–206.
- 90 Croft M, Bradley LM, Swain SL. Naïve versus memory CD4 T cell response to antigen. Memory cells are less dependent on accessory cell costimulation and can respond to many antigen-presenting cell types including resting B cells. *J Immunol* 1994; **152**:2675–85.
- 91 Bruno L, Kirberg J, Von Boehmer H. On the cellular basis of immunological T-cell memory. *Immunity* 1995; **2**:37–43.
- 92 Garcia S, DiSanto J, Stockinger B. Following the development of a CD4 T cell response *in vivo*: from activation to memory formation. *Immunity* 1999; **11**:163–71.
- 93 Veiga-Fernandes H, Walter U, Bourgeois C, McLean A, Rocha B. Response of naïve and memory CD8<sup>+</sup> T cells to antigen stimulation *in vivo*. *Nature Immunol* 2000; **1**:47–53.
- 94 Agarwal S, Rao A. Modulation of chromatin structure regulates cytokine gene expression during T cell differentiation. *Immunity* 1998; **9**:765–75.
- 95 Fitzpatrick DR, Shirley KM, McDonald LE, Bielefeldt-Ohmann H, Kay GF, Kelso A. Distinct methylation of the interferon gamma (IFN- $\gamma$ ) and interleukin 3 (IL-3) genes in newly activated primary CD8<sup>+</sup> T lymphocytes: regional IFN- $\gamma$  promoter demethylation in CD44<sup>high</sup> CD8<sup>+</sup> T cells. *J Exp Med* 1998; **188**:103–17.
- 96 Farber DL, Acuto O, Bottomly K. Differential T cell receptor-mediated signalling in naïve and memory T cells. *Eur J Immunol* 1997; **27**:2094–101.

- 97 Hussain SF, Anderson CF, Farber DL. Differential SLP-76 expression and TCR-mediated signaling in effector and memory CD4 T cells. *J Immunol* 2002; **168**:1557–65.
- 98 Dianzani U, Luqman M, Rojo J, Yagi J, Baron JL, Woods A, Janeway CA, Bottomly K. Molecular associations on the T cell surface correlate with immunological memory. *Eur J Immunol* 1990; **20**:2249–57.
- 99 Leitenberg D, Novak TJ, Farber D, Smith BR, Bottomly K. The extracellular domain of CD45 controls association with the CD4-T cell receptor complex and the response to antigen-specific stimulation. *J Exp Med* 1996; **183**:249–59.
- 100 McHeyzer-Williams MG, Davis MM. Antigen-specific development of primary and memory T cells in vivo. *Science* 1995; **268**:106–11.
- 101 Maryanski JL, Jongeneel CV, Bucher P, Casanova JL, Walker PR. Single-cell PCR analysis of TCR repertoires selected by antigen in vivo: a high magnitude CD8 response is comprised of very few clones. *Immunity* 1996; **4**:47–55.
- 102 Zheng B, Han S, Zhu Q, Goldsby R, Kelsoe G. Alternative pathways for the selection of antigen-specific peripheral T cells. *Nature* 1996; **384**:263–6.
- 103 Busch DH, Pilip I, Pamer EG. Evolution of a complex T cell receptor repertoire during primary and recall bacterial infection. *J Exp Med* 1998; **188**:61–70.
- 104 Busch DH, Pamer EG. T cell affinity maturation by selective expansion during infection. *J Exp Med* 1999; **189**:701–9.
- 105 Rees W, Bender J, Teague TK, Kedl RM, Crawford F, Marrack P, Kappler J. An inverse relationship between T cell receptor affinity and antigen dose during CD4<sup>+</sup> T cell responses in vivo and in vitro. *Proc Natl Acad Sci USA* 1999; **96**:9781–6.
- 106 Savage PA, Boniface JJ, Davis MM. A kinetic basis for T cell receptor repertoire selection during an immune response. *Immunity* 1999; **10**:485–92.
- 107 Fasso M, Anandasabapathy N, Crawford F, Kappler J, Fathman CG, Ridgway WM. T cell receptor (TCR)-mediated repertoire selection and loss of TCR  $\nu\beta$  diversity during the initiation of a CD4<sup>+</sup> T cell response in vivo. *J Exp Med* 2000; **192**:1719–30.
- 108 Kedl RM, Rees WA, Hildeman DA, Schaefer B, Mitchell T, Kappler J, Marrack P. T cells compete for access to antigen-bearing antigen-presenting cells. *J Exp Med* 2000; **192**:1105–13.
- 109 van Bergen J, Kooy Y, Koning F. CD4-independent T cells impair TCR triggering of CD4-dependent T cells: a putative mechanism for T cell affinity maturation. *Eur J Immunol* 2001; **31**:646–52.
- 110 Kedl RM, Schaefer BC, Kappler JW, Marrack P. T cells down-modulate peptide-MHC complexes on APCs *in vivo*. *Nature Immunol* 2002; **3**:27–32.
- 111 Sprent J. Circulating T and B lymphocytes of the mouse. I. Migratory properties. *Cell Immunol* 1973; **7**:10–39.
- 112 Potsch C, Vohringer D, Pircher H. Distinct migration patterns of naive and effector CD8 T cells in the spleen: correlation with CCR7 receptor expression and chemokine reactivity. *Eur J Immunol* 1999; **29**:3562–70.
- 113 Price PW, Cerny J. Characterization of CD4<sup>+</sup> T cells in mouse bone marrow. I. Increased activated/memory phenotype and altered TCR  $\nu\beta$  repertoire. *Eur J Immunol* 1999; **29**:1051–6.
- 114 Masopust D, Vezys V, Marzo AL, Lefrancois L. Preferential localization of effector memory cells in nonlymphoid tissue. *Science* 2001; **291**:2413–6.
- 115 Hogan RJ, Usherwood EJ, Zhong W, Roberts AD, Dutton RW, Harmsen AG, Woodland DL. Activated antigen-specific CD8<sup>+</sup> T cells persist in the lungs following recovery from respiratory virus infections. *J Immunol* 2001; **166**:1813–22.
- 116 Hogan RJ, Zhong W, Usherwood EJ, Cookenham T, Roberts AD, Woodland DL. Protection from respiratory virus infections can be mediated by antigen-specific CD4<sup>+</sup> T cells that persist in the lungs. *J Exp Med* 2001; **193**:981–6.
- 117 Picker LJ, Terstappen LW, Rott LS, Streeter PR, Stein H, Butcher EC. Differential expression of homing-associated adhesion molecules by T cell subsets in man. *J Immunol* 1990; **145**:3247–55.
- 118 Horgan KJ, Luce GE, Tanaka Y, Schweighoffer T, Shimizu Y, Sharrow SO, Shaw S. Differential expression of VLA-alpha 4 and VLA-beta 1 discriminates multiple subsets of CD4<sup>+</sup>CD45RO<sup>+</sup> 'memory' T cells. *J Immunol* 1992; **149**:4082–7.
- 119 Mackay CR, Marston WL, Dudley L, Spertini O, Tedder TF, Hein WR. Tissue-specific migration pathways by phenotypically distinct subpopulations of memory T cells. *Eur J Immunol* 1992; **22**:887–95.
- 120 Schweighoffer T, Tanaka Y, Tidswell M, Erle DJ, Horgan KJ, Luce GEG, Lazarovits AI, Buck D, Shaw S. Selective expression of integrin  $\alpha 4\beta 7$  on a subset of human CD4<sup>+</sup> memory T cells with hallmarks of gut-trophism. *J Immunol* 1993; **151**:717–29.
- 121 Austrup F, Vestweber D, Borges E, Lohning M, Brquer R, Herz U, Renz H, Hallmann R, Scheffold A, Radbruch A, Hamann A. P- and E-selectin mediate recruitment of T-helper-1 but not T-helper-2 cells into inflamed tissues. *Nature* 1997; **385**:81–3.
- 122 Campbell DJ, Butcher EC. Rapid acquisition of tissue-specific homing phenotypes by CD4<sup>+</sup> T cells activated in cutaneous or mucosal lymphoid tissues. *J Exp Med* 2002; **195**:135–41.
- 123 Gallichan WS, Rosenthal KL. Long-lived cytotoxic T lymphocyte memory in mucosal tissues after mucosal but not systemic immunisation. *J Exp Med* 1996; **184**:1879–90.
- 124 Banchereau J, Briere F, Caux C, Davoust J, Lebecque S, Liu YJ, Pulendran B, Palucka K. Immunobiology of dendritic cells. *Annu Rev Immunol* 2000; **18**:767–811.
- 125 Everett NB, Caffrey RW, Rieke WO. Recirculation of lymphocytes. *Ann NY Acad Sci* 1964; **113**:887–97.
- 126 Everett NB, Tyler RW. Lymphopoiesis in the thymus and other tissues: functional implications. *Int Rev Cytol* 1967; **22**:205–37.
- 127 Sprent J, Basten A. Circulating T and B lymphocytes of the mouse II. Lifespan. *Cell Immunol* 1973; **7**:40–59.
- 128 Claesson MH, Ropke C, Hougén HP. Distribution of short-lived and long-lived small lymphocytes in the lymphomyeloid tissues of germ-free NMRI mice. *Scand J Immunol* 1974; **3**:597–604.
- 129 Ropke C. Renewal rates of murine T-lymphocyte subsets. *Cell Immunol* 1990; **128**:185–97.
- 130 Hellerstein M, Hanley MB, Cesar D, Siler S, Papageorgopoulos C, Wieder E, Schmidt D, Hoh R, Neese R, Macallan D, Deeks S, McCune JM. Directly measured kinetics of circulating T lymphocytes in normal and HIV-1 infected humans. *Nature Med* 1999; **5**:83–9.
- 131 Tough DF, Sprent J. Turnover of naive- and memory-phenotype T cells. *J Exp Med* 1994; **179**:1127–35.
- 132 Mohri H, Bonhoeffer S, Monard S, Perelson AS, Ho DD. Rapid turnover of T lymphocytes in SIV-infected Rhesus macaques. *Science* 1998; **279**:1223–7.
- 133 Michie CA, McLean A, Alcock A, Beverley PCL. Lifespan of human lymphocyte subsets defined by CD45 isoforms. *Nature* 1992; **360**:264–5.

- 134 McCune JM, Hanley MB, Cesar D, Halvorsen R, Hoh R, Schmidt D, Wieder E, Deeks S, Siler S, Neese R, Hellerstein M. Factors influencing T-cell turnover in HIV-1-seropositive patients. *J Clin Invest* 2000; **105**:R1–R8.
- 135 Bruno L, von Boehmer H, Kirberg J. Cell division in the compartment of naive and memory T lymphocytes. *Eur J Immunol* 1996; **26**:3179–84.
- 136 Zimmerman C, Brduscha-Riem K, Blaser C, Zinkernagel RM, Pircher H. Visualization, characterization, and turnover of CD8+ memory T cells in virus-infected hosts. *J Exp Med* 1996; **183**:1367–75.
- 137 Takeda S, Rodewald H-R, Arakawa H, Bluethmann H, Shimizu T. MHC class II molecules are not required for survival of newly generated CD4+ T cells, but affect their long-term life span. *Immunity* 1996; **5**:217–28.
- 138 Brocker T. Survival of mature CD4 T lymphocytes is dependent on major histocompatibility complex class II-expressing dendritic cells. *J Exp Med* 1997; **186**:1223–32.
- 139 Kirberg J, Berns A, von Boehmer H. Peripheral T cell survival requires continual ligation of the T cell receptor to major histocompatibility complex-encoded molecules. *J Exp Med* 1997; **186**:1269–75.
- 140 Rooke R, Waltzinger C, Benoist C, Mathis D. Targeted complementation of MHC class II deficiency by intrathymic delivery of recombinant adenoviruses. *Immunity* 1997; **7**:123–34.
- 141 Tanchot C, Lemonnier FA, Perarnau B, Freitas AA, Rocha B. Differential requirements for survival and proliferation of CD8 naive or memory T cells. *Science* 1997; **276**:2057–62.
- 142 Netic D, Vukmanovic S. MHC class I is required for peripheral accumulation of CD8+ thymic emigrants. *J Immunol* 1998; **160**:3705–12.
- 143 Dorfman JR, Stefanova I, Yasutomo K, Germain RN. CD4+ T cell survival is not directly linked to self-MHC-induced TCR signalling. *Nature Immunol* 2000; **1**:329–35.
- 144 Clarke SRM, Rudensky AY. Survival and homeostatic proliferation of naive peripheral CD4+ T cells in the absence of self peptide: MHC complexes. *J Immunol* 2000; **165**:2458–64.
- 145 Witherden D, van Oers N, Waltzinger C, Weiss A, Benoist C, Mathis D. Tetracycline-controllable selection of CD4+ T cells: half-life and survival signals in the absence of major histocompatibility complex class II molecules. *J Exp Med* 2000; **191**:355–64.
- 146 Goldrath AW, Bevan MJ. Selecting and maintaining a diverse T-cell repertoire. *Nature* 1999; **402**:255–62.
- 147 Surh C, Sprent J. Homeostatic T cell proliferation: how far can T cells be activated by self-ligands? *J Exp Med* 2000; **192**:F9–F14.
- 148 Bender J, Mitchell T, Kappler J, Marrack P. CD4+ T cell division in irradiated mice requires peptides distinct from those responsible for thymic selection. *J Exp Med* 1999; **190**:367–73.
- 149 Ahmed R, Gray D. Immunological memory and protective immunity: Understanding their relation. *Science* 1996; **272**:54–60.
- 150 Zinkernagel RM, Bachmann MF, Kundig TM, Oehen S, Pircher H, Hengartner H. On immunological memory. *Annu Rev Immunol* 1996; **14**:333–67.
- 151 Kundig TM, Bachmann MF, Oehen S, Hoffman UW, Simard J, Kalberer CP, Pircher H, Ohashi PS, Hengartner HRM. On the role of antigen in maintaining cytotoxic T-cell memory. *Proc Natl Acad Sci USA* 1996; **93**:9716–23.
- 152 Bachmann MF, Kundig TM, Hengartner H, Zinkernagel RM. Protection against immunopathological consequences of a viral infection by activated but not resting cytotoxic T cells: T cell memory without 'memory T cells'? *Proc Natl Acad Sci USA* 1997; **94**:640–5.
- 153 Zinkernagel RM. What is missing in immunology to understand immunity? *Nature Immunol* 2000; **1**:181–5.
- 154 Hou S, Hyland L, Ryan KW, Portner A, Doherty PC. Virus-specific CD8+ T cell memory determined by clonal burst size. *Nature* 1994; **369**:652–4.
- 155 Mullbacher A. The long-term maintenance of cytotoxic T cell memory does not require persistence of antigen. *J Exp Med* 1994; **179**:317–21.
- 156 Markiewicz MA, Girao C, Opferman JT, Sun J, Hu Q, Agulnik AA, Bishop CE, Thompson CB, Ashton-Rickardt PG. Long-term T cell memory requires the surface expression of self-peptide/major histocompatibility complex molecules. *Proc Natl Acad Sci USA* 1998; **95**:3065–70.
- 157 Swain SL, Hu H, Huston G. Class II-independent generation of CD4 memory T cells from effectors. *Science* 1999; **286**:1381–3.
- 158 Hu H, Huston G, Duso D, Lepak N, Roman E, Swain SL. CD4+ T cell effectors can become memory cells with high efficiency and without further division. *Nature Immunol* 2001; **2**:705–10.
- 159 Kassiotis G, Garcia S, Simpson E, Stockinger B. Impairment of immunological memory in the absence of MHC despite survival of memory T cells. *Nature Immunol* 2002; **3**:244–50.
- 160 Tan JT, Dudl E, LeRoy E, Murray R, Sprent J, Weinberg KI, Surh CD. IL-7 is critical for homeostatic proliferation and survival of naive T cells. *Proc Natl Acad Sci USA* 2001; **98**:8732–7.
- 161 Vivien L, Benoist C, Mathis D. T lymphocytes need IL-7 but not IL-4 or IL-6 to survive in vivo. *Int Immunol* 2001; **13**:763–8.
- 162 Teague TK, Marrack P, Kappler JW, Vella AT. Interleukin-6 rescues naive T cells from apoptosis. *J Immunol* 1997; **158**:5791–6.
- 163 Vella A, Teague TK, Ihle J, Kappler J, Marrack P. Interleukin 4 (IL-4) or IL-7 prevent the death of resting T cells. Stat6 is probably not required for the effect of IL-4. *J Exp Med* 1997; **186**:325–30.
- 164 Tough DF, Borrow P, Sprent J. Induction of bystander T cell proliferation by viruses and type I interferon in vivo. *Science* 1996; **272**:1947–50.
- 165 Tough DF, Sun S, Sprent J. T cell stimulation in vivo by lipopolysaccharide (LPS). *J Exp Med* 1997; **185**:2089–94.
- 166 Tough DF, Zhang X, Sprent J. An IFN- $\gamma$ -dependent pathway controls stimulation of memory phenotype CD8+ T cell turnover in vivo by IL-12, IL-18, and IFN- $\gamma$ . *J Immunol* 2001; **166**:6007–11.
- 167 Ku CC, Murakami M, Sakamoto A, Kappler J, Marrack P. Control of homeostasis of CD8+ memory T cells by opposing cytokines. *Science* 2000; **288**:675–8.
- 168 Lodolce JP, Boone DL, Chai S, Swain RE, Dassopoulos T, Trettin S, Ma A. IL-15 receptor maintains lymphoid homeostasis by supporting lymphocyte homing and proliferation. *Immunity* 1998; **9**:669–76.
- 169 Kennedy MK, Glaccum M, Brown SN, Butz EA, Viney JL, Embers M, Matsuki N, Charrier K, Sedger L, Willis CR, Brasel K, Morrissey PJ, Stocking K, Schuh JCL, Joyce S, Peschon JJ. Reversible defects in natural killer and memory CD8 T cell lineages in interleukin 15-deficient mice. *J Exp Med* 2000; **191**:771–80.
- 170 Nishimura H, Yajima T, Naiki Y, Tsunobuchi H, Umemura M, Itano K, Matsuguchi T, Suzuki M, Ohashi P, Yoshikai Y. Differential roles of interleukin 15 mRNA

- isoforms generated by alternative splicing in immune responses in vivo. *J Exp Med* 2000; **191**:157–69.
- 171 Marks-Konczalik J, Dubois S, Losi JM, Sabzevari H, Yamada N, Feigenbaum L, Waldmann RA, Tagaya Y. IL-2-induced activation-induced cell death is inhibited in IL-15 transgenic mice. *Proc Natl Acad Sci USA* 2000; **97**:445–50.
- 172 Eberl G, Brawand P, MacDonald HR. Selective bystander proliferation of memory CD4<sup>+</sup> and CD8<sup>+</sup> T cells upon NK T or T cell activation. *J Immunol* 2000; **165**:4305–11.
- 173 Geginat J, Sallusto F, Lanzavecchia A. Cytokine-driven proliferation and differentiation of human naive, central memory, and effector memory CD4<sup>+</sup> T cells. *J Exp Med* 2001; **194**:1711–9.
- 174 Cerwenka A, Morgan TM, Dutton RW. Naive, effector, and memory CD8 T cells in protection against pulmonary influenza virus infection: homing properties rather than initial frequencies are crucial. *J Immunol* 1999; **163**:5535–43.